

A group with three real irreducible characters: answering a question of Moretó and Navarro

Mark L. Lewis

*Department of Mathematical Sciences, Kent State University
Kent, Ohio 44242*

E-mail: lewis@math.kent.edu

October 2, 2008

Abstract

In this paper, we construct a group with three real irreducible characters whose Sylow 2-subgroup is an iterated central extension of a Suzuki 2-group. This answers a question raised by Moretó and Navarro who asked whether such a group exists.

MSC primary: 20C15

1 Introduction

In their note [5], Moretó and Navarro study groups G that have 3 real-valued irreducible characters. They prove that G is solvable, and it either has a cyclic Sylow 2-subgroup or it has a normal Sylow 2-subgroup that is either homocyclic, quaternion of order 8, or an iterated central extension of a Suzuki 2-group whose center is an elementary abelian 2-group.

Recall that P is a Suzuki 2-group if P is a nonabelian 2-group, P has more than one involution, and P has a solvable group of automorphisms that transitively permutes the involutions in P . Suzuki 2-groups are studied extensively in Section VIII.7 of [1]. The group P is an iterated central extension of a Suzuki 2-group, if $P/Z_n(P)$ is a Suzuki 2-group for some $n \geq 2$ (where $Z_n(P)$ is the n th term in the upper central series for P).

Now, they present the following examples in that paper. First, S_3 is an example of such a group with a nonnormal, cyclic Sylow 2-subgroup. Also, $\text{SL}(2, 3)$ is such a group with the quaternions as a normal Sylow 2-subgroup. They also produce an example with a normal Sylow 2-subgroup that is a Suzuki 2-group (not of type A). Finally, they include an example with a normal Sylow 2-subgroup that is a central extension of a Suzuki 2-group.

Moretó and Navarro did not have an example where the Sylow 2-subgroup P had $P/Z_n(P)$ is a Suzuki 2-group for $n \geq 2$ where $Z_n(P)$ is the n th term in the upper central series, and they ask whether the word “iterated” can be removed from the statement of their theorem. In this note, we will show that this cannot be done by presenting an example of a group G with three real valued characters whose Sylow 2-subgroup P satisfies $P/Z_2(P)$

is a Suzuki 2-group. At this time, we have not determined whether or not there can be any examples with $n \geq 3$.

We will first discuss the construction used to obtain our group. We then prove that this group has exactly three real irreducible characters. In an appendix, we include the MAGMA code for PC-group arising from our construction, and we discuss the character table of this group.

2 The construction

We now work to construct a group G with 3 real valued irreducible characters whose Sylow 2-subgroup is an iterated Suzuki 2-group. We begin with a group first introduced by Isaacs in [3]. Let F be the field of order 8.

We first establish some facts regarding F . We define the *trace map* $Tr : F \rightarrow Z_2$ by $Tr(\alpha) = \alpha + \alpha^2 + \alpha^4$ for all $\alpha \in F$. It is not difficult to see that Tr is an additive homomorphism. Also, one can show that $\ker(Tr) = \{\alpha + \alpha^2 \mid \alpha \in F\}$. Note that $|\ker(Tr)| = 4$, and in fact, $F = Z_2 \oplus \ker(Tr)$. If we fix an element $b \in F$, we can define a map $f : F \rightarrow F$ by $f(a) = ab^4 + a^4b$. It is easy to show that f is an additive homomorphism whose kernel is $\{0, b\}$.

Consider the skew polynomial ring $F\{X\}$ where we define the multiplication of the indeterminate X with elements of F by $X\alpha = \alpha^2X$ for all $\alpha \in F$. We then define the ring R by $R = F\{X\}/(X^5)$, and we let x be the image of X in R . It is not difficult to see that the Jacobson radical J of R has the form $\{\alpha_1x + \alpha_2x^2 + \alpha_3x^3 + \alpha_4x^4 \mid \alpha_i \in F\}$, and we let $Q = 1 + J$ as a subgroup of the group of units of R . Observe that $|Q| = 2^{12}$. We set $Q^i = 1 + J^i$ for $i = 1, \dots, 5$. We know from [3] that Q^i is a subgroup of Q . Also, $Q = Q^1$ and $Q^5 = 1$.

Let $C = F^\times$, the multiplicative group of F , and we see that C embeds in the group of units of R by $c \in C$ maps to the constant polynomial $c1$. It follows that C acts by conjugation on Q by $(1 + \alpha_1x + \alpha_2x^2 + \alpha_3x^3 + \alpha_4x^4)^c = 1 + \alpha_1cx + \alpha_2c^3x^2 + \alpha_3x^3 + \alpha_4cx^4$. Observe that C acts irreducibly on Q/Q^2 , Q^2/Q^3 , and $Q^4/1$, but C centralizes Q^3/Q^4 . Let $\mathcal{G}$ be the Galois group of F over Z_2 , and we have an action of $\mathcal{G}$ on QC by acting coordinate-wise on the elements of Q and on C as on F . Notice that $C\mathcal{G}$ is a Frobenius group of order 21.

The group Q has a $C\mathcal{G}$ -invariant subgroup P of index 2. To see this, observe that $Q^3/Q^4 = \mathbf{C}_{Q^3/Q^4}(\mathcal{G}) \times [Q^3, \mathcal{G}]Q^4/Q^4$ via Fitting's lemma. We have $\mathbf{C}_{Q^3/Q^4}(\mathcal{G}) = \{1 + \alpha x^3 + Q^4 \mid \alpha \in Z_2\}$, and we set $X/Q^4 = [Q^3, \mathcal{G}]Q^4/Q^4 = \{1 + \alpha x^3 + Q^4 \mid \alpha \in \ker(Tr)\}$. We note that $|Q^3 : X| = 2$. One can also show that $[Q, Q^2]Q^4/Q^4 = X/Q^4$ (this takes some work), and so, $X = [Q, Q^2]Q^4$. Also, we see that

$$X = \{1 + \alpha_3x^3 + \alpha_4x^4 \mid \alpha_3 \in \ker(Tr), \alpha_4 \in F\}.$$

Now, C stabilizes X , so C acts on Q^2/X . Note that Q^2/X is central in Q/X , and thus, Q^2/X is abelian. By Fitting's lemma, we have $Q^2/X = \mathbf{C}_{Q^2/X}(C) \times [Q^2, C]X/X$. We see that $\mathbf{C}_{Q^2/X}(C) = Q^3/X$. Let $Y = [Q^2, C]X$, and observe that $|Q^2 : Y| = |Q^3 : X| = 2$. We see that $Q^2/Y = \mathbf{C}_{Q/Y}(C)$. Also, we claim that

$$Y = \{1 + \alpha_2x^2 + \alpha_3x^3 + \alpha_4x^4 \mid \alpha_2, \alpha_4 \in F, \alpha_3 \in \ker(Tr)\}.$$

Using an argument similar to the one found in Theorem 4.5 of [4], we can prove that $Y = Q'Q^4$. In particular, Q/Y is abelian, so $Q/Y = \mathbf{C}_{Q/Y}(C) \times [Q, C]Y/Y$. We take $P = [Q, C]Y$. Observe that $P' = Q' = Y$. We set $P^i = P \cap Q^i$, so $P^1 = P = [Q, C]Y$, $P^2 = Y$, $P^3 = X$, and $P^4 = Q^4$. We have $Z_1(P) = P^4$, $Z_2(P) = P^3$, and $Z_3(P) = P^2$. In particular, $P/Z_2(P) = P/P^3 \cong Q/Q^3$ which is isomorphic to the Suzuki 2-group of type A and order 64. This proves that P is an iterated Suzuki 2-group.

Our group G is the semidirect product of $C\mathcal{G}$ acting on P .

3 The theorem

Following [2], we will say that $g \in G$ is a *real* element if g is conjugate to g^{-1} . A conjugacy class of G is called *real* if it consists of real elements. In Problem 6.13 of [2], one is asked to prove that the number of real irreducible characters of G equals the number of real classes in G .

Theorem 1. *The group G has exactly three real irreducible characters.*

Proof. We now work to prove that G has exactly 3 irreducible characters. Now G/P is the Frobenius group of order 21, and as such, the principal character is the only real valued character in $\text{Irr}(G/P)$. Let N/P be the Frobenius kernel of G/P , and observe that $|G : N| = 3$. We note that N/P^3 is one of the examples in [5] where $\text{Irr}(N/P^3)$ has two real characters. It is not difficult to show that this will imply that $\text{Irr}(G/P^3)$ has two real characters. Working as in [4], one can show that the nonprincipal characters in $\text{Irr}(P^3/P^4)$ are fully ramified with respect to P/P^3 , so the characters in $\text{Irr}(P/P^4 \mid P^3/P^4)$ are real-valued. If $\theta \in \text{Irr}(P/P^4)$, then the stabilizer of θ in G is N . Now, θ has a unique real extension to N . Hence, θ^G has a unique real irreducible constituent. Note that $\mathcal{G}$ acts transitively on $\text{Irr}(P^3/P^4) - \{1\}$, so, $\text{Irr}(G/P^4 \mid P^3/P^4) = \text{Irr}(G \mid \theta)$. We conclude that $\text{Irr}(G/P^4)$ has three real characters.

We will show that G has 3 real irreducible characters by showing that G has 3 real classes. Notice that G/P^4 has 3 real classes. Also, it is not difficult to see that the nonidentity elements of P^2/P^4 consist of involutions. Notice that the elements of $P^3/P^4 \setminus P^4/P^4$ and $P^2/P^4 \setminus P^3/P^4$ must be in different conjugacy classes. It follows that all of the real elements of G/P^4 lie in P^2/P^4 . Since a real element of G gives rise to a real element of G/P^4 , it follows that the all real elements of G lie in P^2 . In addition, notice since $|G : P|$ is odd, that any element of P^2 that is real in G must in fact be real in P .

A typical element of $a \in Q$ has the form $a = 1 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$ with $a_i \in F$. It is not difficult to see that

$$a^2 = 1 + a_1^3x^2 + (a_1a_2^2 + a_1^4a_2)x^3 + (a_1a_3^2 + a_2^5 + a_1a_3)x^4.$$

It follows that $a^2 = 1$ if and only if $a_1 = a_2 = 0$. Hence, all of the involutions in Q lie in Q^3 , and all of the nonidentity elements of Q^3 are involutions. This implies that all of the involutions in P lie in P^3 , and all of the nonidentity elements of P^3 are involutions. We know that C acts transitively on $P^4 \setminus \{1\}$. So we have that $\{1\}$ and $P^4 \setminus \{1\}$ are two real conjugacy classes in G .

Consider an element $b \in P^2$. We know that $b = 1 + b_2x^2 + b_3x^3 + b_4x^4$ with $b_2, b_4 \in F$ and $b_3 \in \ker(\text{Tr})$. Let a be any element of Q as in the previous paragraph. Computing, we see that

$$ab = 1 + a_1x + (a_2 + b_2)x^2 + (a_3 + b_3 + a_1b_2^2)x^3 + (a_4 + b_4 + a_1b_3^2 + a_2b_2^4)x^4.$$

and

$$ba = 1 + a_1x + (a_2 + b_2)x^2 + (a_3 + b_3 + b_2a_1^4)x^3 + (a_4 + b_4 + b_3a_1 + b_2a_2^4)x^4.$$

It follows that $ab = ba$ if and only if (1) $a_1b_2^2 = b_2a_1^4$ and (2) $a_1b_3^2 + a_2b_2^4 = b_3a_1 + b_2a_2^4$. If $b \in P^3 \setminus P^4$, then $b_2 = 0$ and $b_3 \neq 0$. We see that $ab = ba$ if and only if $a_1b_3^2 = b_3a_1$. Recall that $b_3 \in \ker(\text{Tr})$, so $b_3 \neq 1$. It follows that $b_3 \neq b_3^3$, and hence, $a_1 = 0$. We conclude $\mathbf{C}_Q(b) = Q^2$, and so, $\mathbf{C}_P(b) = P^2$. We then have that $\mathbf{C}_G(b) = P^2C$. It follows that $|G: \mathbf{C}_G(b)| = 8 \cdot 3 = 24$, and so the conjugacy class containing b has size 24. On the other hand, $|P^3 \setminus P^4| = |P^3| - |P^4| = 32 - 8 = 24$. Thus, $P^3 \setminus P^2$ form the third real conjugacy class of G .

We will now show that these are the only real conjugacy classes in G . As we have seen, P^2 contains all of the real classes in G . Thus, we consider $b \in P^2 \setminus P^3$. This implies that $b_2 \neq 0$. It is not difficult to see that the only possible solutions for a_1 in the equation $a_1b_2^2 = a_1^4b_2$ are $a_1 = 0$ and $a_1 = b_2^5$. If $a_1 = 0$, then equation (2) becomes $a_2b_2^4 = a_2^4b_2$. It is not difficult to see that the only solutions for a_2 in this equation are $a_2 = 0$ and $a_2 = b_2$. On the other hand, if $a_1 = b_2^5$, then equation (2) becomes $b_2^5b_3^2 + a_2b_2^4 = b_3b_2^5 + b_2a_2^4$. This implies that $a_2b_2^4 + a_2^4b_2 = b_2^5(b_3 + b_3^2)$. We have seen that the map $a_2 \mapsto a_2b_2^4 + a_2^4b_2$ is an additive homomorphism with a kernel of size 2. This implies that our equation has 2 solutions for a_2 . It is not difficult to see that $P^3 \leq \mathbf{C}_P(b)$. It follows that $|P: \mathbf{C}_P(b)| = 16$. Since $\mathbf{C}_P(b)$ will contain a conjugate of $\mathcal{G}$, we deduce that $|G: \mathbf{C}_G(b)| = 16 \cdot 7 = 112$.

Note that $|P^2 \setminus P^3| = |P^2| - |P^3| = 256 - 32 = 224$. Therefore, $P^2 \setminus P^3$ consists of 2 conjugacy classes of G . This implies that either every element of $P^2 \setminus P^3$ is conjugate to its inverse, or none of the elements in $P^2 \setminus P^3$ are conjugate to their inverse. Thus, we will fix $b = 1 + x^2$. It suffices to show that b is not conjugate to b^{-1} in G . As we have seen, it suffices to show that b is not conjugate to b^{-1} in P . We will show that in fact b is not conjugate to b^{-1} in P .

By way of contradiction, we suppose that b is conjugate to b^{-1} in Q . Hence, there is an element $a \in Q$ so $b^a = b^{-1}$. Observe that $b^{-1} = 1 + x^2 + x^4$. As in the previous paragraph, we can write $a = 1 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$. Let $c = b^a$, and observe that $ac = ba$. Observe that $ba = 1 + a_1x + (a_2 + 1)x^2 + (a_3 + a_1^4)x^3 + (a_4 + a_2^4)x^4$. We know that $c \in P^2$, so $c = 1 + c_2x^2 + c_3x^3 + c_4x^4$, and

$$ac = 1 + a_1x + (a_2 + c_2)x^2 + (a_3 + c_3 + a_1c_2^2)x^3 + (a_4 + c_4 + a_1c_3^2 + a_2c_2^4)x^4.$$

This implies that $c_2 = 1$ and $c_3 = a_1^4 + a_1$ and

$$c_4 = a_2^4 + a_1(a_1^4 + a_1)^2 + a_2 = (a_2^4 + a_2) + (a_1^4 + a_1)^2.$$

If $c = b^{-1}$, then matching coefficients, we must have $a_1^4 + a_1 = 0$ and $(a_2^4 + a_2) + (a_1^4 + a_1)^2 = 1$. Combining these two equations, we obtain $a_2^4 + a_2 = 1$. However, $a_2^4 + a_2 \in \ker(\text{Tr})$ and $1 \notin \ker(\text{Tr})$, and so, we have a contradiction. This contradiction implies that b is not conjugate to its inverse. Therefore, G has 3 real classes and hence, 3 real valued irreducible characters. $\square$

A natural extension of this group is to start our construction by starting with the ring $F\{X\}/\langle X^6 \rangle$, and then completing the construction as before. To see that the resulting group has more than three real irreducible characters, we note that

$$\{1 + a_3x^3 + a_4x^4 + a_5x^5 \mid a_3 \in \ker(Tr), a_4, a_5 \in F\}$$

is a characteristic subgroup of the resulting group. It is not difficult to see that this subgroup has all its nonidentity elements are involutions. Also, it is not difficult to see that the involutions form at least 3 conjugacy classes, and so, the group has at least 4 real conjugacy classes.

4 Appendix: Magma code

Next, include code (in Magma) to generate G as PC-group:

```

> g := PolycyclicGroup < s, c, x1, x2, x3, y1, y2, y3, w1, w2, z1, z2, z3 |
> s^3, c^7, c^s = c^4,
> x1^2 = y1, x2^2 = y2, x3^2 = y3, y1^2 = z1, y2^2 = z2, y3^2 = z3,
> w1^2, w2^2, z1^2, z2^2, z3^2,
> x2^x1 = x2 * y2 * y3 * w1 * w2 * z1 * z2 * z3, x3^x1 = x3 * y1 * y2 * w1 * z2,
> x3^x2 = x3 * y1 * w1 * w2 * z1 * z3,
> y2^x1 = y2 * w1, y3^x1 = y3 * w2,
> y2^y1 = y2 * z3, y3^y1 = y3 * z2 * z3, y3^y2 = y3 * z1 * z2,
> y1^x2 = y1 * w1 * w2 * z1 * z2 * z3, y3^x2 = y3 * w1 * z1,
> y1^x3 = y1 * w1 * z1 * z2, y2^x3 = y2 * w1 * w2 * z2,
> w1^x1 = w1 * z3, w2^x1 = w2 * z2 * z3, w1^x2 = w1 * z1 * z2, w2^x2 = w2 * z1 * z2 * z3,
> w1^x3 = w1 * z2 * z3, w2^x3 = w2 * z1 * z3,
> x1^c = x1 * x2 * x3 * w1 * z1, x2^c = x1 * x3 * y1 * y3 * w1 * z3, x3^c = x1,
> y1^c = y1 * y2 * z1 * z2, y2^c = y2 * y3 * z2 * z3, y3^c = y1,
> w1^c = w1 * z2, w2^c = w2 * z1 * z2 * z3,
> z1^c = z1 * z2 * z3, z2^c = z1 * z3, z3^c = z1,
> x1^s = x1 * z3, x2^s = x2 * x3 * y2 * y3 * w1 * w2 * z1 * z3, x3^s = x2,
> y2^s = y1 * y2 * y3, y3^s = y2, w1^s = w1 * w2, w2^s = w1, z2^s = z2 * z3, z3^s = z2 >;

```

Using Magma, we computed the character table of the above group. This group has a character table with 33 irreducible characters. Given this size, we have not tried to reproduce this character table in this note. The character degrees of G are $\{1, 3, 7, 14, 24, 56\}$. There are three real irreducible characters. One each of degrees 1, 7, and 24.

References

- [1] B. Huppert and N. Blackburn, “Finite Groups II,” Springer-Verlag, Berlin, 1982.
- [2] I. M. Isaacs, “Character Theory of Finite Groups,” Academic Press, San Diego, California, 1976.

- [3] I. M. Isaacs, Coprime group actions fixing all nonlinear irreducible characters, *Canad. J. Math.*, **41** (1989), 68-82.
- [4] M. L. Lewis, A family of groups containing a nonabelian fully ramified section, *J. Algebra*, **186** (1996), 578-596.
- [5] A. Moretó and G. Navarro, Groups with three real valued irreducible characters, *Israel J. Math.*, **163** (2008), 85-92.